

FERMION MASS HIERARCHY FROM THE SOFT WALL

Antonio Delgado¹ and **David Diego**²

*Department of Physics, 225 Nieuwland Science Hall, U. of Notre Dame,
Notre Dame, IN 46556-5670, USA.*

Abstract

We develop a 5d model for ElectroWeak physics based on a non compact warped extra dimension of finite length, known as the soft wall scenario, where all the dynamical degrees of freedom propagate in the 5d bulk. We solve the equations of motion and find the allowed spectra, showing that the mass of the lightest fermionic mode behaves as a power law of the effective 4d Yukawa coupling constant, with the exponent being the corresponding fermionic 5d bulk mass. Precisely this non universal behavior allows us to reproduce the hierarchy between the Standard Model (SM) fermion masses (from neutrinos to the top quark) with non-hierarchy fermionic bulk masses.

¹antonio.delgado@nd.edu

²david.diego@nd.edu

1 Introduction

The existence of warped extra dimensions has been a source of prolific investigations for the last 10 years because it was shown that it could provide with an alternative explanation of the hierarchy between the Planck and the ElectroWeak scales [1, 2]. The pioneer work by Randall and Sundrum [1] considers a slice of a five dimensional anti de Sitter space spanning between two 3-branes, the UV one accounts for the gravity (Planckian) scale and the IR brane is chosen to have a scale of the order of the ElectroWeak breaking by the exponential suppression of the AdS metric. Another aspect of this set-up that has attracted a lot of attention is its connection to the AdS/CFT duality [3] which states that this 5d theory is equivalent to a 4d CFT coupled to gravity and the interpretation of the IR scale is that it corresponds to the breaking of that conformal symmetry and the appearance of a mass gap and resonances [4].

More recently, an alternative scenario with a (non compact) warped extra dimension has been proposed [5, 6, 7], the so-called soft wall as opposed to the hard wall termination of the Randall-Sundrum model. The motivation for the new scenario, as pointed out in reference [5], comes from the fact that a suitable soft wall configuration may lead to linear Kaluza-Klein (KK) excitations ($m_n \sim n$), unlike the quadratic excitations found in the hard wall models, and hence it may provide a description of the linear spectrum of mesons in QCD [8]. This is what has been called the AdS/QCD approach.

In the soft wall case the (effective) metric is not AdS, instead the warped factor decays faster enough to make the extra dimension of finite length and in this sense a virtual IR boundary is located at infinity but with a finite distance from the UV one and the departure from the AdS behavior is associated to a smooth VEV acquired by some dilaton field, however, since no real IR boundary exists all the matter/radiation content propagate in the 5d bulk, thus the suppression due to the metric is universal (and is absorbed through the wave function normalization). The possible suppression on the lowest modes comes from the binding of the Schrödinger-like potentials governing the profiles of the wave functions [6, 7].

In this work we try to reproduce the hierarchy of the fermion masses of the Standard Model embedding it on a soft wall background. It is worth remarking, nevertheless, that our main concern here is to reproduce the correct order of magnitude for the masses, a more realistic study taking into account all the families and all the ElectroWeak constraints would need to incorporate the CKM matrices, which we leave for future investigations. In any case, the incorporation of the mixing angles would not change the main results we are presenting. Finally, since the topic is relatively new and relatively absent in the literature we think a detailed derivation, although overlapping previous publications, is worthwhile.

The paper is structured as follows: In section 2 we present the model together with the notations and conventions. Section 3 is devoted to solve the equations of motion for the

fermion, Higgs and gauge sectors. Section 4 applies the previous results to the hierarchy of the fermion masses and section 5 contains the conclusions. Finally, at the end of the paper, we have included appendices developing some technical aspects of particular calculations.

2 The Model

Before presenting the model we will set up the notation used: Capital letters from the beginning of the alphabet ($A, B, \dots$) represent (local) Lorentz 5d indices while those from the middle and end of the alphabet ($M, N, R, \dots$) stand for general coordinate 5d indices. Finally, greek letters take into account 4d Lorentz indices.

The 5d Lorentz representation is spanned by the gamma matrices $\gamma^A = (\gamma^\mu, -i\gamma^5)$, with γ^μ the usual 4d gamma matrices and

$$\gamma^5 = \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix}.$$

Our model is defined in a five dimensional manifold with (UV) boundary, $\Sigma = R^4 \times \mathcal{I}$, with $\mathcal{I} = [z_0, \infty)$ and $z_0 > 0$, assuming also $z_0 \ll (M_W)^{-1}$ being M_W the ElectroWeak scale. The background metric is AdS, that is

$$g = f^2(z) \eta_5, \quad (1)$$

where $f(z) = k/z$, k being the radius of curvature and as a matter of fact we will take $z_0 \sim k$. In addition η_5 stands for the 5d Minkowski metric with mostly negative signature, namely $\eta_{5AB} = \text{diagonal}(+1, -1, -1, -1, -1)$ and z represents the fifth coordinate. The propagating degrees of freedom consist of an $SU(2)_L$ doublet of 5d Dirac fermions: Ψ_L^i and two singlets Ψ_R^a , an $SU(2)_L$ doublet of complex scalars, H^i , and a gauge field, A_M , lying in the adjoint representation ¹ of $SU(2)_L \times U(1)_Y$, where the lightest modes of the above fields are to be identified with the SM fermions, Higgs and ElectroWeak gauge fields, respectively. For the 5d fields, the bulk-brane action is²

$$\begin{aligned} S = \int_{\Sigma} \sqrt{g} e^{-\phi} & \left[\frac{i}{2} \bar{\Psi}_L^i \gamma^M \mathcal{D}_M^L \Psi_L^i + \frac{i}{2} \bar{\Psi}_R^a \gamma^M \mathcal{D}_M^R \Psi_R^a + M \bar{\Psi}_L^i \Psi_L^i + M \bar{\Psi}_R^a \Psi_R^a \right. \\ & + (\lambda_1 \epsilon_{ij} \bar{\Psi}_L^i \Psi_R^1 H^{*j} + \lambda_2 H^i \bar{\Psi}_L^i \Psi_R^2 + \text{h.c.}) \\ & \left. + g^{MN} (D_M H)^\dagger D_N H - m_h^2 H^\dagger H + \frac{1}{4g_5^2} g^{MR} g^{NS} \text{tr} \{F_{MN} F_{RS}\} \right] \\ & - \int_{\partial\Sigma} \sqrt{g} e^{-\phi} \left[\lambda_0 k^2 (|H|^2 - v_0^2)^2 - e_5^5 \frac{1}{2} (\bar{\Psi}_L^i \Psi_L^i - \bar{\Psi}_R^a \Psi_R^a) \right], \quad (2) \end{aligned}$$

¹To avoid important quantum corrections to the ρ parameter the gauge group should be $SU(2)_L \times SU(2)_R \times U(1)_Y$ [6, 7] although for the sake of simplicity we take the usual gauge group. For the calculations of the spectra this technicality does not affect.

²The presence of the (5,5) component of the inverse vielbein, e_5^5 , in the brane action is not a problem because the (effective) background metric itself breaks the isometry group to 4d Lorentz-Poincaré.

where $F_{MN} = \partial_{[M} A_{N]} + [A_M, A_N]$ and $A_M = g_5 A_M^s T_s$ being the group generators verifying $[T_s, T_t] = if_{st}^r T_r$ and g_5 the (5d) gauge coupling constant, $\lambda^{1,2}$ are the 5d Yukawa coupling constants, M is a (constant) bulk mass and $m_h^2 = \frac{a(a-4)}{k^2} - 2a\frac{\mu^2}{k^2} z^2$ with a, λ_0 being dimensionless constants whereas v_0 has dimension of (energy) $^{3/2}$. ϵ_{ij} is the Levi-Civita tensor and ϕ is the dilaton field. In addition

$$D_M = \partial_M + iA_M, \quad (3)$$

$$\mathcal{D}_M^L = D_M + E_M^C \Gamma_{CB}^A \eta_{AD} \Sigma^{BD}, \quad (4)$$

$$\mathcal{D}_M^R = \partial_M + E_M^C \Gamma_{CB}^A \eta_{AD} \Sigma^{BD}, \quad (5)$$

are gauge and Lorentz covariant derivatives, where E_M^A is the vielbein field, verifying $g_{MN} = E_M^A E_N^B \eta_{AB}$. The gamma matrices depend on the spacetime coordinates according to $\gamma^M = e_A^M \gamma^A$ with e_A^M being the inverse of the vielbein field verifying $g_{MN} e_A^M e_B^N = \eta_{AB}$, i.e. they are the (local coordinate) components of orthonormal vector fields ($e_A \equiv e_A^M \partial_M$). Furthermore, $\Sigma^{AB} = \frac{1}{8} [\gamma^A, \gamma^B]$ are the Lorentz group generators in the representation

$$\text{diagonal} \left(SL(2, \mathbb{C}) \times SL(2, \mathbb{C})^{\dagger -1} \right),$$

and the symbols Γ_{BC}^A verify $\nabla_{e_B} e_C = \Gamma_{BC}^A e_A$ with ∇ the Levi-Civita connection compatible with g . The relation with the Christoffel symbols, Γ_{MN}^R , is then $\Gamma_{BC}^A = e_B^M E_N^A \partial_M e_C^N + e_B^M e_C^N E_R^A \Gamma_{MN}^R$. In the case of the metric (1) the Lorentz covariant derivative is

$$\mathcal{D}_\mu^R = \partial_\mu + \frac{i}{2} \gamma^5 \gamma_\mu \frac{f'}{f}, \quad (6)$$

$$\mathcal{D}_5^R = \partial_5. \quad (7)$$

For the dilaton field we will assume a z -dependent VEV. The particular bulk-brane gravitational dynamic leading to that profile for the dilaton is beyond the scope of the present work, although in Ref. [7] an underlying gravity model leading to a soft wall configuration can be found. Here we will take $\langle \phi \rangle = \mu^2 z^2$ with μ some mass scale. As for the Higgs bulk mass is concerned, its z dependence can be understood as a consequence of the coupling to the gravity sector. We will also not address the question of the different scales in the problem but supposed that their stability is granted by an underlying UV theory.

Finally, we have taken the same (z -independent) bulk mass for Left- and Right-handed fermions for the sake of simplicity at solving the equations of motion. Different masses, as it can be seen in the next section, substantially increase the difficulty as it is not possible to decouple easily the first order fermionic equations of motion.

3 Solving the equations of motion

We will start by finding the VEV of the Higgs following the derivation found in Ref. [7].

3.1 Higgs VEV

Consider the purely scalar sector³ out of the action (2) and assume for the moment $H(x, z) \equiv \mathfrak{H}(z)$, then, the minimal action principle yields the equations of motion

$$\mathfrak{H}'' + \left(3\frac{f'}{f} - \phi'\right) \mathfrak{H}' - f^2 m_h^2 \mathfrak{H} = 0, \quad (8)$$

together with the boundary conditions

$$\mathfrak{H}' - 2\lambda_0 k^2 (\mathfrak{H}^2 - v_0^2) \mathfrak{H}|_{z_0} = 0. \quad (9)$$

Redefining the scalar as $\mathfrak{H} = z^a \mathcal{H}$ we find the equation

$$\mathcal{H}'' + \left(\frac{2a-3}{z} - 2\mu^2 z\right) \mathcal{H}' = 0, \quad (10)$$

and thus the general solution is

$$\mathfrak{H} = z^a \left(c + \tilde{c} \int_{z_0}^z d\zeta \zeta^{3-2a} e^{\mu^2 \zeta^2} \right), \quad (11)$$

although the normalizable solution is obtained with $\tilde{c} = 0$ and hence the boundary conditions set the constraint

$$c [a - 2\lambda_0 k^2 (c^2 z_0^{2a} - v_0^2) z_0] = 0, \quad (12)$$

whose solution is

$$\begin{cases} c = 0 \\ c^2 = \left(\frac{1}{z_0}\right)^{2a} \left(\frac{a}{2z_0 \lambda_0 k^2} + v_0^2\right) \end{cases} \quad (13)$$

Now if we compute the energy density per unit 4d volume [9] for the solutions to the system (8)-(9) we find

$$\begin{aligned} \mathcal{E}_{\mathfrak{H}} &= \int_{\Sigma} \sqrt{g} e^{-\phi} (-g^{55} \partial_5 \mathfrak{H} \partial_5 \mathfrak{H} + m_h^2 \mathfrak{H}^2) + \int_{\partial\Sigma} \sqrt{g} e^{-\phi} \lambda_0 k^2 (\mathfrak{H}^2 - v_0^2)^2 \\ &= \int_{\Sigma} \sqrt{g} e^{-\phi} \mathfrak{H} [g^{55} \nabla_5 \partial_5 \mathfrak{H} - g^{55} \partial_5 \phi \partial_5 \mathfrak{H} + m_h^2 \mathfrak{H}] \\ &\quad + \int_{\partial\Sigma} \sqrt{g} e^{-\phi} [\mathfrak{H} \mathfrak{H}' + \lambda_0 k^2 (\mathfrak{H}^2 - v_0^2)^2] \\ &= \lambda_0 k^2 \int_{\partial\Sigma} \sqrt{g} e^{-\phi} (v_0^4 - \mathfrak{H}^4). \end{aligned} \quad (14)$$

The energy density per unit 4d volume is then decreasing around the trivial solution, $\mathfrak{H} = 0$, for $\lambda_0 > 0$ and therefore in this case the non trivial solution is the absolute minimum.

Now that the Higgs profile has been set, we then proceed by solving the equations of motion for the fermion sector according to three polynomial cases for the Higgs VEV.

³We consider the Higgs doublet in the unitary gauge, that is $H = \begin{pmatrix} 0 \\ \mathfrak{H} \end{pmatrix}$ with $\mathfrak{H} \in \mathbb{R}$.

3.2 Fermionic sector

Applying the variational principle to the fermionic sector⁴ of the action (2) one finds a bulk piece whose vanishing induces the following equations of motion

$$ie_A^M \gamma^A \mathcal{D}_M^R \Psi_L^i - \frac{1}{2} e_A^5 \gamma^A \phi' \Psi_L^i + M \Psi_L^i + \lambda_1 \epsilon_{ij} \mathfrak{H}^j \Psi_R^1 + \lambda_2 \mathfrak{H}^i \Psi_R^2 = 0, \quad (15)$$

$$ie_A^M \gamma^A \mathcal{D}_M^R \Psi_R^1 - \frac{1}{2} e_A^5 \gamma^A \phi' \Psi_R^1 + M \Psi_R^1 + \lambda_1 \epsilon_{ij} \Psi_L^i \mathfrak{H}^j = 0, \quad (16)$$

$$ie_A^M \gamma^A \mathcal{D}_M^R \Psi_R^2 - \frac{1}{2} e_A^5 \gamma^A \phi' \Psi_R^2 + M \Psi_R^2 + \lambda_2 \mathfrak{H}^i \Psi_L^i = 0 \quad (17)$$

while the boundary variation yields the constraint

$$e_5^5 (\mathbf{1} - \sigma_3 \otimes \gamma^5) \begin{pmatrix} \Psi_L^i \\ \Psi_R^a \end{pmatrix}_{z_0} = 0. \quad (18)$$

where σ_3 is the z -Pauli matrix acting on (L, R) space. If we refer Ψ_L^i and Ψ_R^a as simply Ψ_L and Ψ_R , respectively, the equations of motion and the boundary conditions adopt the generic form

$$ie_A^M \gamma^A \mathcal{D}_M^R \Psi_L - \frac{1}{2} e_A^5 \gamma^A \phi' \Psi_L + M \Psi_L + \lambda \mathfrak{H}(z) \Psi_R, \quad (19)$$

$$ie_A^M \gamma^A \mathcal{D}_M^R \Psi_R - \frac{1}{2} e_A^5 \gamma^A \phi' \Psi_R + M \Psi_R + \lambda \mathfrak{H}(z) \Psi_L = 0, \quad (20)$$

$$e_5^5 (\mathbf{1} - \sigma_3 \otimes \gamma^5) \begin{pmatrix} \Psi_L \\ \Psi_R \end{pmatrix}_{z_0} = 0. \quad (21)$$

and upon the redefinitions⁵

$$\Psi_{L,R} = \frac{1}{\sqrt{2}} f^{-2} e^{\frac{1}{2} \phi} (\psi_+ \mp \psi_-), \quad (22)$$

we find

$$i \not{\partial} \psi_{\pm} + \gamma^5 \partial_5 \psi_{\pm} + (M \pm m_D) f \psi_{\pm} = 0, \quad (23)$$

with $m_D = \lambda \mathfrak{H}(z)$, while the boundary conditions take the form

$$(\mathbf{1} - \sigma_1 \otimes \gamma^5) \begin{pmatrix} \psi_+ \\ \psi_- \end{pmatrix} = 0. \quad (24)$$

Considering now an orbifold-like decomposition⁶, that is

$$\psi_{\pm} = \begin{pmatrix} h_{\pm}(z) \xi_{\pm}(x^{\mu}) \\ g_{\pm}(z) \bar{\chi}_{\pm}(x^{\mu}) \end{pmatrix}, \quad (25)$$

⁴Considering the Lorentz covariant derivative only.

⁵With different L and R fermionic bulk masses one can not decouple the system with a global rotation.

⁶Notice that the normalization of the kinetic term in these variables is simply $\int_{z_0}^{\infty} |h|^2$.

where $\xi_{\pm}, \chi_{\pm}$ are Weyl spinors verifying $i\bar{\sigma}^{\mu}\partial_{\mu}\xi_{\pm} = m\bar{\chi}_{\pm}$, m being the physical eigenmass, the first order coupled differential equations (23) can be written as

$$h'_{\pm} + (M \pm m_D) f h_{\pm} = -m g_{\pm}, \quad (26)$$

$$g'_{\pm} - (M \pm m_D) f g_{\pm} = m h_{\pm}, \quad (27)$$

and differentiating once more they transform into two decoupled 2nd order differential equations

$$h''_{\pm} - \left[f^2 (M \pm m_D)^2 - f' (M \pm m_D) \mp f m'_D - m^2 \right] h_{\pm} = 0, \quad (28)$$

$$g''_{\pm} - \left[f^2 (M \pm m_D)^2 + f' (M \pm m_D) \pm f m'_D - m^2 \right] g_{\pm} = 0. \quad (29)$$

In addition the boundary conditions (24), in terms of the new variables, translate into⁷

$$(\mathbf{1} - \sigma_1) \begin{pmatrix} h_+ \\ h_- \end{pmatrix} = (\mathbf{1} + \sigma_1) \begin{pmatrix} g_+ \\ g_- \end{pmatrix} = 0. \quad (30)$$

In the following we explicitly solve these equations for three polynomial behaviors of the Higgs VEV.

- Constant profile ($a = 0$): $\mathfrak{H}(z) = v_0$

In this case the equations take the form

$$\frac{d}{d\theta} h_{\pm} + \frac{\alpha_{\pm}}{\theta} h_{\pm} = -g_{\pm}, \quad (31)$$

$$\frac{d}{d\theta} g_{\pm} - \frac{\alpha_{\pm}}{\theta} g_{\pm} = h_{\pm}, \quad (32)$$

$$\frac{d^2}{d\theta^2} h_{\pm} - \frac{\alpha_{\pm}(\alpha_{\pm} + 1)}{\theta^2} h_{\pm} + h_{\pm} = 0, \quad (33)$$

$$\frac{d^2}{d\theta^2} g_{\pm} - \frac{\alpha_{\pm}(\alpha_{\pm} - 1)}{\theta^2} g_{\pm} + g_{\pm} = 0, \quad (34)$$

with $\alpha_{\pm} = Mk \pm \lambda v_0 k$ and $\theta = mz$. Redefining the fuctions as $h_{\pm} = \sqrt{\theta} \mathfrak{h}_{\pm}$ and $g_{\pm} = \sqrt{\theta} \mathfrak{g}_{\pm}$ we find the Bessel equations

$$\theta \frac{d}{d\theta} \left(\theta \frac{d}{d\theta} \mathfrak{h}_{\pm} \right) + \left[\theta^2 - \left(\alpha_{\pm} + \frac{1}{2} \right)^2 \right] \mathfrak{h}_{\pm} = 0, \quad (35)$$

$$\theta \frac{d}{d\theta} \left(\theta \frac{d}{d\theta} \mathfrak{g}_{\pm} \right) + \left[\theta^2 - \left(\alpha_{\pm} - \frac{1}{2} \right)^2 \right] \mathfrak{g}_{\pm} = 0, \quad (36)$$

⁷Eq. (24) implies $\chi_+ = \kappa \chi_-$ with κ a constant (and an analogous relation for $\xi_{\pm}$) however by redefining the Weyl spinors one can absorb the constants.

and, accordingly, the solutions are

$$h_{\pm} = \sqrt{\theta} \left[H_{\pm}^J J \left(\alpha_{\pm} + \frac{1}{2}, \theta \right) + H_{\pm}^Y Y \left(\alpha_{\pm} + \frac{1}{2}, \theta \right) \right], \quad (37)$$

$$g_{\pm} = \sqrt{\theta} \left[G_{\pm}^J J \left(\alpha_{\pm} - \frac{1}{2}, \theta \right) + G_{\pm}^Y Y \left(\alpha_{\pm} - \frac{1}{2}, \theta \right) \right], \quad (38)$$

for $\alpha_{\pm} \pm \frac{1}{2}$ non-integer. One can easily check that

$$\left(\theta \frac{d}{d\theta} + \alpha_{\pm} \right) \left[\sqrt{\theta} J \left(\frac{1}{2} + \alpha_{\pm}, \theta \right) \right] = \theta \sqrt{\theta} J \left(-\frac{1}{2} + \alpha_{\pm}, \theta \right), \quad (39)$$

$$\left(\theta \frac{d}{d\theta} + \alpha_{\pm} \right) \left[\sqrt{\theta} Y \left(\frac{1}{2} + \alpha_{\pm}, \theta \right) \right] = \theta \sqrt{\theta} Y \left(-\frac{1}{2} + \alpha_{\pm}, \theta \right), \quad (40)$$

and hence the first order constraints (31)-(32) impose

$$G_{\pm}^J = -H_{\pm}^J, \quad (41)$$

$$G_{\pm}^Y = -H_{\pm}^Y. \quad (42)$$

Since we have enough degrees of freedom to solve the boundary conditions (30), no further restriction on the mass eigenvalue is found. Thus, the spectrum is continuum and there is no mass gap.

- Linear profile ($a = 1$): $\mathfrak{H}(z) = \frac{\tilde{\mu}}{\sqrt{k}} \frac{z}{k}$ with $\tilde{\mu} = \frac{1}{z_0} \sqrt{\frac{k}{2z_0\lambda_0} + v_0^2 k^3}$.

The equations of motion are now

$$\frac{d}{d\Theta} h_{\pm} + \left(\frac{Mk}{\Theta} \pm \frac{\lambda}{2\sqrt{k}} \frac{\tilde{\mu}}{\gamma} \right) h_{\pm} = -\frac{m}{2\gamma} g_{\pm}, \quad (43)$$

$$\frac{d}{d\Theta} g_{\pm} - \left(\frac{Mk}{\Theta} \pm \frac{\lambda}{2\sqrt{k}} \frac{\tilde{\mu}}{\gamma} \right) g_{\pm} = \frac{m}{2\gamma} h_{\pm}, \quad (44)$$

$$\frac{d^2}{d\Theta^2} h_{\pm} - \left[\frac{Mk(Mk+1)}{\Theta^2} \pm \frac{M\lambda\tilde{\mu}\sqrt{k}}{\gamma\Theta} + \frac{1}{4} \right] h_{\pm} = 0, \quad (45)$$

$$\frac{d^2}{d\Theta^2} g_{\pm} - \left[\frac{Mk(Mk-1)}{\Theta^2} \pm \frac{M\lambda\tilde{\mu}\sqrt{k}}{\gamma\Theta} + \frac{1}{4} \right] g_{\pm} = 0, \quad (46)$$

with $\Theta = 2\gamma z$ and $\gamma = \sqrt{\frac{\lambda^2}{k} \tilde{\mu}^2 - m^2}$. Then with the redefinitions

$$h_{\pm} = e^{-\Theta/2} \Theta^{1+Mk} \mathfrak{h}_{\pm}, \quad g_{\pm} = e^{-\Theta/2} \Theta^{1-Mk} \mathfrak{g}_{\pm}, \quad (47)$$

Eqs. (45)-(46) turn into the following confluent hypergeometric equations (alsol known as Kummer equations)

$$\Theta \frac{d^2}{d\Theta^2} \mathfrak{h}_{\pm} + (2 + 2\alpha - \Theta) \frac{d}{d\Theta} \mathfrak{h}_{\pm} - (1 + \alpha \pm \beta) \mathfrak{h}_{\pm} = 0, \quad (48)$$

$$\Theta \frac{d^2}{d\Theta^2} \mathfrak{g}_{\pm} + (2 - 2\alpha - \Theta) \frac{d}{d\Theta} \mathfrak{g}_{\pm} - (1 - \alpha \pm \beta) \mathfrak{g}_{\pm} = 0, \quad (49)$$

where $\alpha = Mk$ and $\beta = \frac{M\lambda\tilde{\mu}\sqrt{k}}{\gamma}$. For $2 \pm 2\alpha \notin \mathbb{Z}$ the general solution is (see appendix A)

$$\mathfrak{h}_{\pm} = H_{\pm} {}_1F_1(1 + \alpha \pm \beta, 2 + 2\alpha, \Theta) + H'_{\pm} U(1 + \alpha \pm \beta, 2 + 2\alpha, \Theta), \quad (50)$$

$$\mathfrak{g}_{\pm} = G_{\pm} {}_1F_1(1 - \alpha \pm \beta, 2 - 2\alpha, \Theta) + G'_{\pm} U(1 - \alpha \pm \beta, 2 - 2\alpha, \Theta). \quad (51)$$

Furthermore, one easily show that

$$\left[\frac{d}{d\Theta} + \left(\frac{\alpha}{\Theta} \pm \frac{\beta}{2\alpha} \right) \right] \left\{ \Theta^{1+\alpha} e^{-\Theta/2} {}_1F_1(1 + \alpha \pm \beta, 2 + 2\alpha, \Theta) \right\} = \\ (1 + 2\alpha) \Theta^{\alpha} e^{-\Theta/2} {}_1F_1(-\alpha \pm \beta, -2\alpha, \Theta), \quad (52)$$

$$\left[\frac{d}{d\Theta} + \left(\frac{\alpha}{\Theta} \pm \frac{\beta}{2\alpha} \right) \right] \left\{ \Theta^{1+\alpha} e^{-\Theta/2} U(1 + \alpha \pm \beta, 2 + 2\alpha, \Theta) \right\} = \\ \frac{-\alpha \pm \beta}{2\alpha} \Theta^{\alpha} e^{-\Theta/2} U(-\alpha \pm \beta, -2\alpha, \Theta), \quad (53)$$

and so for $M \rightarrow -M$ and $h \rightarrow g$, thus the general solution to the coupled system (43)-(44) is given by

$$h_{\pm} = \Theta^{1+\alpha} e^{-\Theta/2} [f_{\pm} {}_1F_1(1 + \alpha \pm \beta, 2 + 2\alpha, \Theta) \\ + u_{\pm} U(1 + \alpha \pm \beta, 2 + 2\alpha, \Theta)], \quad (54)$$

$$g_{\pm} = \Theta^{\alpha} e^{-\Theta/2} \left[-f_{\pm} \frac{2\gamma(1 + 2\alpha)}{m} {}_1F_1(-\alpha \pm \beta, -2\alpha, \Theta) \right. \\ \left. + u_{\pm} \frac{\gamma(\alpha \mp \beta)}{m\alpha} U(-\alpha \pm \beta, -2\alpha, \Theta) \right], \quad (55)$$

As far as the spectrum is concerned we find two distinct regions:

1. $m^2 > \frac{\lambda^2 \tilde{\mu}^2}{k}$, i.e. $\Theta = i|\Theta|$ (continuum spectrum)

In this case none of the above linearly independent solutions are normalizable and analogously to the previous case the boundary conditions at z_0 are satisfied without further restrictions on the physical mass.

2. $m^2 < \frac{\lambda^2 \tilde{\mu}^2}{k}$ (discrete spectrum)

For real Θ only the U -type solution is normalizable since ${}_1F_1(\alpha, \beta, \Theta) \sim \Theta^{\alpha-\beta} e^{\Theta}$ while $U(\alpha, \beta, \Theta) \sim \Theta^{-\alpha}$ as $\Theta \rightarrow \infty$. We then set $f_{\pm} = 0$ and thus a non trivial solution to the boundary conditions (30) requires the vanishing of the function

$$(\alpha + \beta) U(1 + \alpha + \beta, 2 + 2\alpha, \Theta_0) U(-\alpha - \beta, -2\alpha, \Theta_0) \\ + (\alpha - \beta) U(1 + \alpha - \beta, 2 + 2\alpha, \Theta_0) U(-\alpha + \beta, -2\alpha, \Theta_0),$$

with $\Theta_0 = 2\gamma z_0 \ll 1$.

Therefore the linear case predicts a finite set of eigenmasses followed by a continuum of eigenvalues. However the mass restriction does not allow a light mode.

- Quadratic profile ($a = 2$): $\mathfrak{H}(z) = \frac{\bar{\mu}}{\sqrt{k}} \frac{z^2}{k^2}$ with $\bar{\mu} = \left(\frac{\sqrt{k}}{z_0}\right)^2 \sqrt{\frac{k}{z_0 \lambda_0} + v_0^2 k^3}$.

In this last case, the (first order) equations of motion turn into

$$\frac{d}{d\zeta} h_{\pm} + \frac{1}{\zeta} (\alpha \pm \zeta) h_{\pm} = -\delta \frac{1}{\sqrt{\zeta}} g_{\pm}, \quad (56)$$

$$\frac{d}{d\zeta} g_{\pm} - \frac{1}{\zeta} (\alpha \pm \zeta) g_{\pm} = \delta \frac{1}{\sqrt{\zeta}} h_{\pm}, \quad (57)$$

with $\zeta = \left(\frac{z}{z_0}\right)^2 \zeta_0$, $\alpha = Mk$, $\delta = \frac{mz_0}{\sqrt{\zeta_0}}$ and $\zeta_0 = \frac{\lambda}{\sqrt{k}} \sqrt{\frac{k}{z_0 \lambda_0} + v_0^2 k^3}$. Redefining the functions as

$$\begin{aligned} h_+ &= \zeta^{-\frac{\alpha}{2}} e^{-\frac{1}{2}\zeta} \mathfrak{h}_+, & g_- &= \zeta^{\frac{\alpha}{2}} e^{-\frac{1}{2}\zeta} \mathfrak{g}_-, \\ h_- &= \zeta^{\frac{1+\alpha}{2}} e^{-\frac{1}{2}\zeta} \mathfrak{h}_-, & g_+ &= \zeta^{\frac{1-\alpha}{2}} e^{-\frac{1}{2}\zeta} \mathfrak{g}_+, \end{aligned}$$

the second order differential equations become the following Kummer's equations

$$\zeta \mathfrak{h}_+'' + \left(\frac{1}{2} - \alpha - \zeta\right) \mathfrak{h}_+' + \frac{\delta^2}{4} \mathfrak{h}_+ = 0, \quad (58)$$

$$\zeta \mathfrak{g}_-'' + \left(\frac{1}{2} + \alpha - \zeta\right) \mathfrak{g}_-' + \frac{\delta^2}{4} \mathfrak{g}_- = 0, \quad (59)$$

$$\zeta \mathfrak{h}_-'' + \left(\frac{3}{2} + \alpha - \zeta\right) \mathfrak{h}_-' - \left(1 - \frac{\delta^2}{4}\right) \mathfrak{h}_- = 0, \quad (60)$$

$$\zeta \mathfrak{g}_+'' + \left(\frac{3}{2} - \alpha - \zeta\right) \mathfrak{g}_+' - \left(1 - \frac{\delta^2}{4}\right) \mathfrak{g}_+ = 0, \quad (61)$$

such that the (normalizable) solutions are

$$h_+ = a_+ \zeta^{-\frac{\alpha}{2}} e^{-\frac{1}{2}\zeta} U\left(-\frac{\delta^2}{4}, \frac{1}{2} - \alpha, \zeta\right), \quad (62)$$

$$h_- = a_- \zeta^{\frac{1+\alpha}{2}} e^{-\frac{1}{2}\zeta} U\left(1 - \frac{\delta^2}{4}, \frac{3}{2} + \alpha, \zeta\right), \quad (63)$$

$$g_+ = b_+ \zeta^{\frac{1-\alpha}{2}} e^{-\frac{1}{2}\zeta} U\left(1 - \frac{\delta^2}{4}, \frac{3}{2} - \alpha, \zeta\right), \quad (64)$$

$$g_- = b_- \zeta^{\frac{\alpha}{2}} e^{-\frac{1}{2}\zeta} U\left(-\frac{\delta^2}{4}, \frac{1}{2} + \alpha, \zeta\right), \quad (65)$$

while the first order constraints impose

$$b_+ = -\frac{1}{2}\delta a_+, \quad a_- = \frac{1}{2}\delta b_-, \quad (66)$$




Figure 1: *Normalized fermionic mass quantization condition as a function of δ for $\zeta_0 = 10^{-3}$ and $\alpha = 3$ (solid line) and $\alpha = 5$ (dashed line).*

then the solutions are given by

$$\psi_+ = \begin{pmatrix} \zeta^{-\frac{\alpha}{2}} e^{-\frac{1}{2}\zeta} U\left(-\frac{\delta^2}{4}, \frac{1}{2} - \alpha, \zeta\right) \xi(x) \\ -\frac{\delta}{2} \zeta^{\frac{1-\alpha}{2}} e^{-\frac{1}{2}\zeta} U\left(1 - \frac{\delta^2}{4}, \frac{3}{2} - \alpha, \zeta\right) \bar{\chi}(x) \end{pmatrix}, \quad (67)$$

$$\psi_- = \begin{pmatrix} \frac{\delta}{2} \zeta^{\frac{1+\alpha}{2}} e^{-\frac{1}{2}\zeta} U\left(1 - \frac{\delta^2}{4}, \frac{3}{2} + \alpha, \zeta\right) \xi(x) \\ \zeta^{\frac{\alpha}{2}} e^{-\frac{1}{2}\zeta} U\left(-\frac{\delta^2}{4}, \frac{1}{2} + \alpha, \zeta\right) \bar{\chi}(x) \end{pmatrix}. \quad (68)$$

Finally, a non trivial solution out of the boundary conditions (30) set the restriction

$$\begin{aligned} & U\left(-\frac{\delta^2}{4}, \frac{1}{2} - \alpha, \zeta_0\right) U\left(-\frac{\delta^2}{4}, \frac{1}{2} + \alpha, \zeta_0\right) \\ & - \frac{1}{4} \delta^2 \zeta_0 U\left(1 - \frac{\delta^2}{4}, \frac{3}{2} - \alpha, \zeta_0\right) U\left(1 - \frac{\delta^2}{4}, \frac{3}{2} + \alpha, \zeta_0\right) = 0. \end{aligned}$$

This condition quantizes the possible physical masses and indeed it presents a mass gap. To see this notice that the confluent hypergeometric function $U(a, b, z)$, as a function of a , oscillates for negative values of this parameter, then we expect to find oscillatory behavior for $\delta \gtrsim 2$. Actually, in Fig. 1 we plot the above mass quantization condition for $\zeta_0 \sim 10^{-3}$ and two different values of α .

Further assuming that $\zeta_0 \ll 1$ (at the end of section 4 we justify this smallness) and for $|\alpha| > 1/2$ we find a light smallest Dirac eigenmass, given by

$$m^2 k^2 \approx 2 \frac{1}{\Gamma(|\alpha| - \frac{1}{2})} (\zeta_0)^{|\alpha| + \frac{1}{2}}. \quad (69)$$

For $|\alpha| < 1/2$ the lowest eigenvalue is not power law suppressed [7]. Indeed the authors of Ref. [7] propose such different behavior as a source of the hierarchy. Nevertheless, as

we shall see, the hierarchy can be induced within the power law behavior. As for the spectrum is concerned, we are interested in a discrete set of KK excitations, hence from now on we will consider a quadratic profile for the Higgs VEV and as pointed out in the previous section, the lightest modes will be identified as the SM fields.

To finish this section we will present the solutions for the bosonic sector.

3.3 Bosonic sector

3.3.1 ElectroWeak equations of motion

The variational principle applied on the gauge piece of the action (2) (taking into account the Higgs VEV) sets the equations of motion

$$\frac{1}{\sqrt{g}} \partial_N \{ \sqrt{g} g^{MR} g^{NS} \partial_{[R} A_{S]} \} - g^{MN} g^{RS} \partial_{[N} A_{R]} \partial_S \phi - g_5^2 \mathfrak{H}^2 g^{MN} A_N = 0, \quad (70)$$

along with the boundary conditions

$$F^{5\mu} = g^{5M} g^{\mu N} \partial_{[M} A_{N]} \big|_{z_0} = 0. \quad (71)$$

Applied to the AdS metric (1) in the gauge $A_5 = 0$ and $\partial_M A^M = 0$ we find the equations of motion

$$\left(-\square + \partial_5^2 + \frac{f'}{f} \partial_5 - \phi' \partial_5 - g_5^2 \mathfrak{H}^2 f^2 \right) A^\mu = 0, \quad (72)$$

$$\partial_5 A^\mu \big|_{z_0} = 0. \quad (73)$$

Decomposing the gauge field as $A^\mu(x, z) = v^\mu(x) g(z)$ with $\square v^\mu = -m^2 v^\mu$ we find

$$g'' + \frac{f'}{f} g' - \phi' g' - (g_5^2 \mathfrak{H}^2 f^2 - m^2) g = 0. \quad (74)$$

Then with the redefinitions

$$\begin{cases} g = f^{-1/2} e^{\frac{1}{2}\phi} e^{-\frac{1}{2}\theta} \theta^{-\frac{1}{4}} \mathfrak{g} \\ \theta = \sqrt{1 + \frac{g_5^2 \bar{\mu}^2}{\mu^4 k^3}} \mu^2 z^2 \end{cases} \quad (75)$$

we find

$$\theta \frac{d^2}{d\theta^2} \mathfrak{g} - \theta \frac{d}{d\theta} \mathfrak{g} + \mathfrak{m}^2 \mathfrak{g} = 0, \quad (76)$$

where $\mathfrak{m} = \frac{m}{2\gamma\mu}$ and $\gamma = \sqrt{1 + \frac{g_5^2 \bar{\mu}^2}{\mu^4 k^3}}$. Finding the general solution to this particular case of Kummer differential equation is somehow involved. Here we simply give the normalizable

solution, although a detailed derivation can be found in appendix A. For the gauge field the m -th mode is

$$A^\mu(x, z) = v^\mu(x) e^{\frac{1}{2}(1-\gamma)\mu^2 z^2} \mathfrak{g}(\gamma\mu^2 z^2), \quad (77)$$

with

$$\mathfrak{g}(\theta) = 1 - \mathfrak{m}^2 \theta \ln \theta {}_1F_1(1 - \mathfrak{m}^2, 2, \theta) + \frac{\mathfrak{m}^2 \theta}{\Gamma(1 - \mathfrak{m}^2)} \mathfrak{u}(\theta), \quad (78)$$

where

$$\mathfrak{u}(\theta) = \sum_{k=0}^{\infty} \frac{\Gamma(1 - \mathfrak{m}^2 + k)}{k! \Gamma(2 + k)} [\Psi_0(1 + k) + \Psi_0(2 + k) - \Psi_0(1 - \mathfrak{m}^2 + k)] \theta^k,$$

and $\Psi_0(x) \equiv \frac{\Gamma'(x)}{\Gamma(x)}$ is the digamma function. The boundary conditions (73) are

$$\frac{1}{2}(1 - \gamma) \mathfrak{g}(\theta_0) + \gamma \mathfrak{g}'(\theta_0) = 0. \quad (79)$$

Assuming then that $\theta_0 \ll 1$ we find for the lowest mode ($\mathfrak{m}^2 \ll 1$)

$$1 - \gamma - 2\gamma\mathfrak{m}^2 [1 - \Psi_0(2) + \ln \theta_0] + \mathcal{O}(\theta_0 \ln \theta_0) = 0, \quad (80)$$

and hence

$$m_W^2 \approx 2\mu^2 \gamma (\gamma - 1) \frac{1}{|\ln \theta_0|}. \quad (81)$$

Notice that γ is always greater or equal than 1, and thus the mass is always real. On the other hand, when the coupling to the Higgs VEV is switched off ($g_5 \rightarrow 0$), which is the case of the photon, the lowest eigenmass is zero. Indeed, for $\gamma = 1$ and $m = 0$ the equations of motion can be written as

$$f e^{-\phi} \partial_5 \left(f e^{-\phi} \partial_5 A^\mu \right) = 0, \quad (82)$$

and together with the boundary condition $f e^{-\phi} \partial_5 A^\mu|_{z_0} = 0$, both impose a constant profile for the massless mode.

3.3.2 Higgs equations of motion

As far as the Higgs is concerned, here we will analyze the perturbations around the background found in section 3.1, that is $H(x, z) = \mathfrak{H}(z) + \tilde{H}(x, z)$, which gives the free equations of motion

$$\square \tilde{H} - \partial_5^2 \tilde{H} - \left(3 \frac{f'}{f} - \phi' \right) \partial_5 \tilde{H} + f^2 m_h^2 \tilde{H} = 0, \quad (83)$$

with $\square \tilde{H} = -m^2 \tilde{H}$, while the boundary variation imposes the vanishing of the expression

$$\partial_5 \tilde{H} - 2\lambda_0 k^2 \left[\left(|\mathfrak{H} + \tilde{H}|^2 - v_0^2 \right) \tilde{H} + |\tilde{H}|^2 \mathfrak{H} + \left(\tilde{H} + \tilde{H}^* \right) \mathfrak{H} \right], \quad (84)$$

at the boundary $z = z_0$. For solving the equations of motion we redefine the the function as $\tilde{H} = f^{-3/2} \tau^{\frac{1}{4}} \mathfrak{h}$ with $\tau = \mu^2 z^2$ and then we find the Kummer equation

$$\tau \frac{d^2}{d\tau^2} \mathfrak{h} + (1 - \tau) \frac{d}{d\tau} \mathfrak{h} + \frac{m^2}{4\mu^2} \mathfrak{h} = 0. \quad (85)$$

Again we refer to appendix A to find the details of the solution and here we simply write down the normalizable one, which turns out to be

$$\begin{aligned} \tilde{H}(x, \tau) = & H(x) \Gamma\left(-\frac{m^2}{4\mu^2}\right) \tau \ln \tau {}_1F_1\left(-\frac{m^2}{4\mu^2}, 1, \tau\right) \\ & - H(x) \tau \sum_{k=0}^{\infty} \frac{\Gamma\left(-\frac{m^2}{4\mu^2} + k\right)}{k! \Gamma(1+k)} \left[2\Psi_0(1+k) - \Psi_0\left(-\frac{m^2}{4\mu^2} + k\right)\right] \tau^k, \end{aligned} \quad (86)$$

or shortly: $\tilde{H}(x, z) \equiv H(x) \eta(\mu^2 z^2)$. Assuming that $\tau_0 \ll 1$ the boundary conditions can be expressed as

$$1 + \ln \tau_0 - 2\Psi_0(1) + \frac{\Gamma'\left(-\frac{m^2}{4\mu^2}\right)}{\Gamma\left(-\frac{m^2}{4\mu^2}\right)} + \mathcal{O}(\tau_0 \ln \tau_0) = 0, \quad (87)$$

and since $\frac{\Gamma'(\epsilon)}{\Gamma(\epsilon)} \sim -\frac{1}{\epsilon}$ for $\epsilon \ll 1$ we find that the lowest Higgs mass is

$$m_H^2 \approx 4\mu^2 \frac{1}{|\ln \tau_0|}. \quad (88)$$

4 Phenomenological predictions on the hierarchy of the SM fermion masses

In this section we will apply the previous results to the fermion masses. In particular we will use them as a possible source for the hierarchy of the masses.

Whenever we have normalizable solutions, a 4d effective action is well defined. Starting from the action (2), by means of partial integration and upon the redefinitions (22) we find the action (in appendix C we give a proof for the orthogonality of the normalizable solutions)

$$\sum_{n \geq 0} \int d^4x \left(i \bar{\psi}_{\mathfrak{f}}^{(n)} \not{\partial} \psi_{\mathfrak{f}}^{(n)} - m_{\mathfrak{f}}^{(n)} \bar{\psi}_{\mathfrak{f}}^{(n)} \psi_{\mathfrak{f}}^{(n)} \right), \quad (89)$$

where⁸

$$\psi_{\mathfrak{f}}^{(n)} \equiv \begin{pmatrix} \psi_{\mathfrak{f}L}^{(n)} \\ \psi_{\mathfrak{f}R}^{(n)} \end{pmatrix},$$

⁸Recall that the boundary conditions (24) set $\xi_+ = \xi_-$ and $\chi_+ = \chi_-$ and thus we are left with a single Dirac fermion.

with f labeling the fermion flavour. In addition we will assume $z_0 \sim k$.

Notice, however, that the values of the 5d couplings are physically meaningless by themselves, instead one should consider the effective 4d couplings obtained upon the integration over the fifth dimension, and for this aim we need to normalize the wave functions. The normalization constants are given by the expressions

$$\begin{aligned} \left(N_L^{(m)}\right)^{-2} &= \frac{k}{2\sqrt{\zeta_0}} \int_{\zeta_0}^{\infty} d\zeta e^{-\zeta} \left[\zeta^{-\frac{1}{2}-\alpha} U^2 \left(-\frac{\delta_m^2}{4}, \frac{1}{2} - \alpha, \zeta \right) \right. \\ &\quad \left. + \frac{\delta_m^2}{4} \zeta^{\frac{1}{2}+\alpha} U^2 \left(1 - \frac{\delta_m^2}{4}, \frac{3}{2} + \alpha, \zeta \right) \right], \end{aligned} \quad (90)$$

$$\begin{aligned} \left(N_R^{(r)}\right)^{-2} &= \frac{k}{2\sqrt{\zeta_0}} \int_{\zeta_0}^{\infty} d\zeta e^{-\zeta} \left[\zeta^{-\frac{1}{2}+\alpha} U^2 \left(-\frac{\delta_r^2}{4}, \frac{1}{2} + \alpha, \zeta \right) \right. \\ &\quad \left. + \frac{\delta_r^2}{4} \zeta^{\frac{1}{2}-\alpha} U^2 \left(1 - \frac{\delta_r^2}{4}, \frac{3}{2} - \alpha, \zeta \right) \right], \end{aligned} \quad (91)$$

for the fermions, while for the Higgs one finds

$$\left(N_H^{(n)}\right)^{-2} = \frac{(\tau_0)^{\frac{3}{2}}}{2\mu} \int_{\tau_0}^{\infty} d\tau e^{-\tau} \tau^{-2} \eta_{(n)}^2(\tau). \quad (92)$$

As for the effective Yukawa coupling is concerned, we can read it off from the action (2)

$$\sqrt{g} e^{-\phi} \lambda H (\bar{\Psi}_L \Psi_R + \bar{\Psi}_R \Psi_L) = \lambda f H (\bar{\psi}_+ \psi_+ - \bar{\psi}_- \psi_-), \quad (93)$$

and therefore we find the coupling

$$Y^{(n,m,r)} = y_{f4d}^{(n,m,r)} H^{(n)}(x) \left(\psi_{fL}^{(m)}(x) \psi_{fR}^{(r)}(x) + \text{h.c.} \right), \quad (94)$$

with

$$\begin{aligned} y_{f4d}^{(n,m,r)} &= \frac{k\lambda\tau_0}{4\zeta_0} N_H^{(n)} N_L^{(m)} N_R^{(r)} \int_{\zeta_0}^{\infty} d\zeta e^{-\zeta} \left(\frac{\tau_0}{\zeta_0} \zeta \right)^{-1} \eta_{(n)} \left(\frac{\tau_0}{\zeta_0} \zeta \right) \\ &\quad \times \left[\delta_r \zeta^{\frac{1}{2}+\alpha} U \left(1 - \frac{\delta_m^2}{4}, \frac{3}{2} + \alpha, \zeta \right) U \left(-\frac{\delta_r^2}{4}, \frac{1}{2} + \alpha, \zeta \right) \right. \\ &\quad \left. - \delta_m \zeta^{\frac{1}{2}-\alpha} U \left(1 - \frac{\delta_r^2}{4}, \frac{3}{2} - \alpha, \zeta \right) U \left(-\frac{\delta_m^2}{4}, \frac{1}{2} - \alpha, \zeta \right) \right], \end{aligned} \quad (95)$$

where, for simplicity, we have omitted the flavor index in the integrals. Thus the effective SM Yukawa coupling would be the one between the lightest modes, which will be referred to as simply $y_{4d}^f \equiv y_{f4d}^{(0,0,0)}$. In appendix B we give an estimation of the previous hypergeometric integrals based on the smallness of the parameters τ_0, ζ_0 and the exponential




Figure 2: $-\log_{10}(mk)$ as a function of α for $k^{-1} \sim 10^{19} \text{ GeV}$ and either $y_{4d} \lesssim 1$ (dashed line) or $y_{4d} \sim 0.1$ (solid line).

suppression of the integrands. According to that approximation we can estimate the order of magnitude of the fermionic physical (Dirac) mass as

$$\left\{ \begin{array}{ll} (m_f k)^2 \sim \frac{2}{\Gamma(|\alpha_f| - \frac{1}{2})} \left[\sqrt{\frac{|\alpha_f| - \frac{3}{2}}{|\alpha_f| - \frac{1}{2}}} \frac{y_{f4d}}{|\ln \tau_0|} \right]^{|\alpha_f| + \frac{1}{2}}, & |\alpha| > \frac{3}{2} \\ m_f k \sim \frac{\sqrt{2}}{\Gamma(\frac{3}{2} - \alpha) [\Psi_0(\frac{3}{2} - \alpha) + \ln(\frac{\tau_0}{c_0})]} y_{f4d}, & \frac{1}{2} < |\alpha| < \frac{3}{2} \end{array} \right. \quad (96)$$

Recall that $\alpha_f = M_f k$, M_f being the 5d fermion bulk mass.

We will be interested, nevertheless, in the case $|\alpha| > \frac{3}{2}$. Indeed this power law dependence on the Yukawa coupling allows us to reproduce the hierarchy between the SM fermion masses with a tiny variation in the 5d bulk masses.

To extract some numbers we will take the scale mass μ associated with the dilaton VEV as $\mu \sim \text{TeV}$, indeed this is needed for the gauge and the Higgs masses, Eqs. (81) and (88), respectively, to be at the ElectroWeak scale, and similarly we will take the 5d VEV of the Higgs as $v_0 \sim \frac{1}{\sqrt{k}} \text{TeV}$ and $|\lambda_0| \lesssim 1$. For the curvature, however, we will consider two extreme situations:

1. Planckian curvature, $k^{-1} \sim 10^{19} \text{ GeV}$.

As we can see in Fig. 2 with order 1 effective Yukawa couplings, $y_{4d} \lesssim 1$, we find the correct order of magnitude for the top quark mass ($m_t k \sim 10^{-17}$) for $|\alpha_t| \simeq 13$ and for the neutrinos masses ($m_\nu k \sim 10^{-28}, 10^{-31}$ -depending on whether they are light, $\sim \text{eV}$, or ultralight $\sim \text{meV}$ -) with $|\alpha_\nu| \simeq 20, 22$, respectively.

Of course, the smallness of the mass scale μ compared to the AdS curvature, $\mu k \sim 10^{-16}$, and the stabilization of the dilaton VEV deserve an explanation that we are not addressing in this work.




Figure 3: $-\log_{10}(mk)$ as a function of α for $k^{-1} \sim 10^4 \text{ GeV}$ and either $y_{4d} \lesssim 1$ (dashed line) or $y_{4d} \sim 0.1$ (solid line).

If we, instead, consider slightly suppressed 4d Yukawa couplings, $y_{4d} \sim 0.1$, the suitable values of α are: $|\alpha_t| \simeq 10$ and $|\alpha_\nu| \simeq 15, 17$.

2. LHC curvature, $k^{-1} \sim 10^4 \text{ GeV}$.

In this case (as Fig. 3 shows) with $\mathcal{O}(1)$ Yukawa couplings we reach the order of magnitude for the top quark ($m_t k \sim 10^{-2}$) with $|\alpha_t| \simeq 5$ while for the neutrinos ($m_\nu k \sim 10^{-13}, 10^{-16}$) we have to consider $|\alpha_\nu| \simeq 18, 21$, respectively.

And as before, taking $y_{4d} \sim 0.1$ we instead find $|\alpha_t| \simeq 2$ and $|\alpha_\nu| \simeq 12, 14$.

Concerning the parameters ζ_0^f, τ_0 , from Eq. (69) we can express ζ_0^f as

$$\zeta_0^f = \left[\frac{m_f^2 k^2}{2} \Gamma\left(\alpha_f - \frac{1}{2}\right) \right]^{\frac{1}{\alpha_f + \frac{1}{2}}},$$

from where we find

$$\left\{ \begin{array}{l} k^{-1} \sim 10^{19} \text{ GeV}, \tau_0 \sim 10^{-32}, \left\{ \begin{array}{ll} y_{4d} \lesssim 1, & \zeta_0^f \sim 10^{-2} \\ y_{4d} \sim 0.1, & \zeta_0^f \sim 10^{-3} \end{array} \right. \\ \\ k^{-1} \sim 10^4 \text{ GeV}, \tau_0 \sim 10^{-2}, \left\{ \begin{array}{ll} y_{4d} \lesssim 1, & \zeta_0^f \sim 10^{-1} \\ y_{4d} \sim 0.1, & \zeta_0^f \sim 10^{-2} \end{array} \right. \end{array} \right.$$

One should compare the previous results with the case of Randall-Sundrum, where the hierarchy of fermions, at least if one forgets about neutrinos, is obtained for $O(1)$ parameters whereas here we need to have $O(10)$ parameters. Although this is not a big difference, the reason why we have such a result is that we are enforcing both the left-handed doublet and right-handed singlet to have the same 5d mass whereas in the case

of RS the prescription taken is $c_L = 1/2 - c_R$. Have we taken a different prescription we would have obtained different results, but we would have lost any possibility of solving the problem analytically. Since the whole point of this paper is just to show how one can generate the hierarchy and not to give a complete model we have preferred to be as simple as possible.

Let us finish this section by saying that all the masses that we generate in this set-up are *Dirac* since they come from a Yukawa coupling of the Higgs with two different fermions. This will imply that neutrinos are Dirac particles and hence there will be no neutrinoless double beta decay. If it is indeed found that neutrinos are Majorana particles one can think of accommodating that fact into this model by adding some localized new sterile neutrinos in the UV brane or some mass term, also localized, for the right handed partner of the neutrino. We leave this also for further investigation.

5 Conclusions

We have developed a model for ElectroWeak physics embedded in five (non compact) dimensions with an exponentially decaying effective metric such that the fifth dimension has finite length. The departure of the metric from the AdS case is associated to the VEV acquired by some dilaton field ϕ for which we have taken $\langle\phi\rangle = \mu^2 z^2$, z representing the fifth coordinate and μ being some mass scale, although the underlying gravity model giving room to this particular aspect has not been addressed in the current work.

In addition, the spacetime has a UV boundary but no IR one. This class of models are commonly known as soft wall models in the literature. A smooth fifth coordinate dependent Higgs VEV may be justified within these models, turning out in a power law behavior. Taking this result into account, we have solved the fermionic equations of motion for three different polynomial behaviors of the Higgs VEV: constant, linear and quadratic, where the latter is the only one binding the solutions enough for them to be normalizable and hence yielding a discrete set of KK excitations.

We have further solved the corresponding equations of motion for the Higgs and ElectroWeak sectors and found that the smallest eigenmasses are of the same order of magnitude, both being proportional to μ and thus we are forced to consider the latter at the TeV scale.

Finally, we have found that (the order of magnitude of) the lightest fermionic masses behave as a power law of the effective 4d Yukawa couplings, where the exponent is the product between the corresponding fermionic 5d bulk mass and the (AdS) radius of curvature (k). This power law precisely allows us to reproduce the Standard Model hierarchy between the fermionic masses with a mild variation for the bulk masses and even order 1 effective Yukawa couplings. In particular with a bulk 5d mass ranging on $\sim (10, 20) \frac{1}{k}$ we achieve the correct order of magnitude for the top quark and the neutrinos, respec-

tively, with a Planckian curvature, $k^{-1} \sim 10^{19}\text{GeV}$. While for a much lower curvature, $k^{-1} \sim 10^4\text{GeV}$ we reproduce the hierarchy of the masses with 5d bulk masses taking values on $\sim (2, 20)^{\frac{1}{k}}$. We would like to stress again, however, that our main concern here has been to end up with the correct order of magnitude for the SM fermion masses. A more realistic study incorporating the CKM matrices and the Electroweak constraints, although the main results presented in the current work would not substantially change, alike the question concerning the dilaton VEV and its stabilization should be addressed in future investigations.

A Solution to the Kummer equation for integer first parameter (b)

The general form of the Kummer differential equation is [10]

$$x y'' + (b - x) y' - a y = 0. \quad (97)$$

For $b \notin \mathbb{Z}$ the two linearly independent solutions are

$$y_1 = {}_1F_1(a, b, x) \equiv \frac{\Gamma(b)}{\Gamma(a)} \sum_{k=0}^{\infty} \frac{\Gamma(a+k)}{k! \Gamma(b+k)} x^k, \quad (98)$$

$$y_2 = x^{1-b} {}_1F_1(1+a-b, 2-b, x). \quad (99)$$

For the confluent hypergeometric function of the first kind, ${}_1F_1(a, b, x)$, we have

$${}_1F_1(a, b, x) \approx x^{a-b} e^x \left(\frac{\Gamma(b)}{\Gamma(a)} + \mathcal{O}\left(\frac{1}{x}\right) \right) + x^{-2a} \left(\frac{(-1)^a}{\Gamma(b-a)} + \mathcal{O}\left(\frac{1}{x}\right) \right),$$

thus a non-exponential asymptotic behavior is found with the linear combination⁹

$$U(a, b, x) = \frac{\Gamma(1-b)}{\Gamma(1+a-b)} y_1 + \frac{\Gamma(b-1)}{\Gamma(a)} y_2. \quad (100)$$

For $b \in \mathbb{Z}$ the solutions y_1, y_2 are no longer independent. Actually one easily checks that

$$\lim_{b \rightarrow n} \frac{\Gamma(a)}{\Gamma(b)} {}_1F_1(a, b, z),$$

does exist for any $n \in \mathbb{Z}$. Indeed, when $b \rightarrow -n$ with n a positive or zero integer one finds

$$\frac{\Gamma(a)}{\Gamma(b)} {}_1F_1(a, b, z) \rightarrow \frac{\Gamma(1+a+n)}{\Gamma(2+n)} z^{1+n} {}_1F_1(1+a+n, 2+n, z),$$

and then the function

$$\begin{aligned} \tilde{U}(a, b, z) &\equiv \frac{\Gamma(a)}{\Gamma(b)} {}_1F_1(a, b, z) - \frac{\Gamma(1+a-b)}{\Gamma(2-b)} z^{1-b} {}_1F_1(1+a-b, 2-b, z) \\ &= \frac{\Gamma(1+a-b) \Gamma(a)}{\Gamma(1-b) \Gamma(b)} U(a, b, z), \end{aligned}$$

vanishes as b approaches any integer value. Thus two linearly independent solutions are given by

$$y_1^n = \lim_{b \rightarrow n} \frac{\Gamma(a)}{\Gamma(b)} {}_1F_1(a, b, z), \quad (101)$$

$$y_2^n = \lim_{b \rightarrow n} \frac{(-1)^n \pi}{\sin \pi b} \tilde{U}(a, b, z) = \partial_b \tilde{U}(a, b, z) \Big|_{b=n}, \quad (102)$$

⁹It behaves as a power law.

where for the last equality we have used the l'Hôpital rule. In addition notice that y_2^n behaves as a power law when $z \rightarrow \infty$ since it is the limit of a function proportional to $U(a, b, z)$.

For the particular cases $b = 0$ and $b = 1$, corresponding to the gauge and the Higgs cases, respectively, one easily shows that

$$\lim_{b \rightarrow 0} \partial_b \frac{\Gamma(a)}{\Gamma(b)} {}_1F_1(a, b, z) = \Gamma(a) - z \sum_{k=0}^{\infty} \frac{\Gamma(1+a+k)}{k! \Gamma(2+k)} \Psi_0(1+k) z^k,$$

$$\begin{aligned} \lim_{b \rightarrow 0} \partial_b z^{1-b} \frac{\Gamma(1+a-b)}{\Gamma(2-b)} {}_1F_1(1+a-b, 2-b, z) = \\ z \sum_{k=0}^{\infty} \frac{\Gamma(1+a+k)}{k! \Gamma(2+k)} [\Psi_0(2+k) - \Psi_0(1+a+k)] z^k \\ - \Gamma(1+a) z \ln z {}_1F_1(1+a, 2, z) \end{aligned}$$

$$\lim_{b \rightarrow 1} \partial_b \frac{\Gamma(a)}{\Gamma(b)} {}_1F_1(a, b, z) = - \sum_{k=0}^{\infty} \frac{\Gamma(a+k)}{k! \Gamma(1+k)} \Psi_0(1+k) z^k,$$

$$\begin{aligned} \lim_{b \rightarrow 1} \partial_b z^{1-b} \frac{\Gamma(1+a-b)}{\Gamma(2-b)} {}_1F_1(1+a-b, 2-b, z) = \\ \sum_{k=0}^{\infty} \frac{\Gamma(a+k)}{k! \Gamma(1+k)} [\Psi_0(1+k) - \Psi_0(a+k)] z^k \\ - \Gamma(a) \ln z {}_1F_1(a, 1, z), \end{aligned}$$

with $\Psi_0(x) \equiv \{\ln[\Gamma(x)]\}'$, the digamma function, and hence we find the (normalizable) solutions presented in subsection 3.3.

B Estimate of the hypergeometric integrals

To estimate the value of the wave function normalizations and the effective 4d Yukawa couplings we will assume $\tau_0, \zeta_0 \ll 1$. Furthermore, since all the integrands are exponentially suppressed we will approximate the hypergeometric functions to the lowest order expansion, disregarding the higher values of the integral variable, that is

$$U(a, b, x) \approx \left[\frac{\Gamma(1-b)}{\Gamma(1-b+a)} + \frac{\Gamma(b-1)}{\Gamma(a)} x^{1-b} \right] (1 + \mathcal{O}(x)). \quad (103)$$

- $|\alpha| > \frac{3}{2}$

For the case of the kinetic normalizations we have

$$(N_R)^{-2} = \frac{k}{2\sqrt{\zeta_0}} \frac{\Gamma^2\left(\alpha - \frac{1}{2}\right)}{\Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)} \left[\frac{\delta^2}{4} + \frac{\Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)}{\Gamma^2\left(-\frac{\delta^2}{4}\right)} \right] \\ \times \int_{\zeta_0}^{\infty} d\zeta e^{-\zeta} \zeta^{\frac{1}{2}-\alpha} \left[1 + \mathcal{O}\left(\zeta^{\alpha-\frac{1}{2}}\right) \right]. \quad (104)$$

The leading order integral can be approximated using the L'Hôpital rule when $\zeta_0 \rightarrow 0$ for which we find

$$\int_{\zeta_0}^{\infty} d\zeta e^{-\zeta} \zeta^{\frac{1}{2}-\alpha} \left[1 + \mathcal{O}\left(\zeta^{\alpha-\frac{1}{2}}\right) \right] \approx \frac{(\zeta_0)^{\frac{3}{2}-\alpha}}{\alpha - \frac{3}{2}} e^{-\zeta_0} \left[1 + \mathcal{O}(\zeta_0)^{\alpha-\frac{3}{2}} \right], \quad (105)$$

and then

$$(N_R)^{-2} \approx \frac{k}{2\sqrt{\zeta_0}} \frac{\delta^2}{4} \frac{\Gamma^2\left(\alpha - \frac{1}{2}\right)}{\Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)} \left[1 + \frac{\delta^2 \Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)}{4 \Gamma^2\left(1 - \frac{\delta^2}{4}\right)} \right] \\ \times \frac{(\zeta_0)^{\frac{3}{2}-\alpha}}{\alpha - \frac{3}{2}} e^{-\zeta_0}. \quad (106)$$

Analogously, for the positive parity spinor ψ_L we find

$$(N_L)^{-2} \approx \frac{k}{2\sqrt{\zeta_0}} \frac{\Gamma^2\left(\alpha + \frac{1}{2}\right)}{\Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)} \left[1 + \frac{\delta^2 \Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)}{4 \Gamma^2\left(1 - \frac{\delta^2}{4}\right)} \right] \\ \times \frac{(\zeta_0)^{\frac{1}{2}-\alpha}}{\alpha - \frac{1}{2}} e^{-\zeta_0}. \quad (107)$$

Notice that $N_L(\alpha) = N_R(-\alpha)$ thus if α were negative the roles of N_L and N_R would simply interchange.

For the Higgs we take $\tau^{-1}\eta(\tau) \approx \Gamma\left(-\frac{m_H^2}{4\mu^2}\right) \ln \tau$ and hence

$$(N_H)^{-2} \approx \frac{(\tau_0)^{\frac{3}{2}}}{2\mu} \Gamma^2\left(-\frac{m_H^2}{4\mu^2}\right) \Gamma''(1), \quad (108)$$

where we have approximated

$$\int_{\tau_0}^{\infty} e^{-\tau} \ln^2 \tau \approx \int_0^{\infty} e^{-\tau} \ln^2 \tau = \Gamma''(1). \quad (109)$$

For the effective Yukawa coupling we start from

$$y_{4d} \approx -\frac{\lambda k \delta \tau_0}{4\zeta_0} N_L N_R N_H \Gamma\left(-\frac{m_H^2}{4\mu}\right) \frac{\Gamma\left(\alpha - \frac{1}{2}\right) \Gamma\left(\alpha + \frac{1}{2}\right)}{\Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)} \\ \times \left[1 + \frac{\delta^2 \Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)}{4 \Gamma^2\left(1 - \frac{\delta^2}{4}\right)} \right] \int_{\zeta_0}^{\infty} d\zeta e^{-\zeta} \zeta^{\frac{1}{2}-\alpha} \ln\left(\frac{\tau_0}{\zeta_0} \zeta\right), \quad (110)$$

where we have taken

$$\begin{aligned}
& \zeta^{\frac{1}{2}+\alpha} U\left(1 - \frac{\delta^2}{4}, \frac{3}{2} + \alpha, \zeta\right) U\left(-\frac{\delta^2}{4}, \frac{1}{2} - \alpha, \zeta\right) \\
& - \zeta^{\frac{1}{2}-\alpha} U\left(1 - \frac{\delta^2}{4}, \frac{3}{2} - \alpha, \zeta\right) U\left(-\frac{\delta^2}{4}, \frac{1}{2} - \alpha, \zeta\right) \\
& \approx -\zeta^{\frac{1}{2}-\alpha} \frac{\Gamma\left(\alpha - \frac{1}{2}\right) \Gamma\left(\alpha + \frac{1}{2}\right)}{\Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)} \left[1 + \frac{\delta^2 \Gamma^2\left(\frac{1}{2} + \alpha - \frac{\delta^2}{4}\right)}{4 \Gamma^2\left(1 - \frac{\delta^2}{4}\right)}\right].
\end{aligned} \tag{111}$$

Using again the L'Hôpital rule we find

$$\int_{\zeta_0}^{\infty} d\zeta e^{-\zeta} \zeta^{\frac{1}{2}-\alpha} \ln\left(\frac{\tau_0}{\zeta_0} \zeta\right) \approx \frac{(\zeta_0)^{\frac{3}{2}-\alpha}}{\alpha - \frac{3}{2}} e^{-\zeta_0} \ln \tau_0, \tag{112}$$

and by plugging the normalization constants in the expression (110) and taking $z_0 \sim k$, the former simplifies to

$$y_{4d} \approx \frac{\sqrt{2} \mu k}{\sqrt{\Gamma''(1) \left(\frac{1}{\lambda_0} + v_0^2 k^3\right)}} \zeta_0 |\ln \tau_0| \sqrt{\frac{\alpha - \frac{1}{2}}{\alpha - \frac{3}{2}}}, \tag{113}$$

where we have used that $\zeta_0 = \frac{\lambda}{\sqrt{k}} \sqrt{\frac{1}{\lambda_0} + v_0^2 k^3}$. In addition, from Eq. (69) we can express ζ_0 as

$$\zeta_0 = \left[\frac{m^2 k^2}{2} \Gamma\left(\alpha - \frac{1}{2}\right) \right]^{\frac{1}{\alpha + \frac{1}{2}}}. \tag{114}$$

Finally, since we have set $\mu \sim v_0 \sqrt{k} \sim \text{TeV}$ we may naturally assume

$$\frac{\sqrt{2} \mu k}{\sqrt{\Gamma''(1) \left(\frac{1}{\lambda_0} + v_0^2 k^3\right)}} \sim \mathcal{O}(1), \tag{115}$$

and then, from (113) one obtains the corresponding expression in (96).

- $\frac{1}{2} < |\alpha| < \frac{3}{2}$

If α were such that $1/2 < |\alpha| < 3/2$ we would instead find

$$\begin{aligned}
& \int_{\zeta_0}^{\infty} e^{-\zeta} \zeta^{\frac{1}{2}-\alpha} \approx \Gamma\left(\frac{3}{2} - \alpha\right), \\
& \int_{\zeta_0}^{\infty} e^{-\zeta} \zeta^{-\frac{1}{2}-\alpha} \approx \frac{(\zeta_0)^{\frac{1}{2}-\alpha}}{\alpha - \frac{1}{2}} e^{-\zeta_0}, \\
& \int_{\zeta_0}^{\infty} e^{-\zeta} \zeta^{\frac{1}{2}-\alpha} \ln\left(\frac{\tau_0}{\zeta_0} \zeta\right) \approx \Gamma\left(\frac{3}{2} - \alpha\right) \left[\Psi_0\left(\frac{3}{2} - \alpha\right) + \ln\left(\frac{\tau_0}{\zeta_0}\right) \right],
\end{aligned}$$

with $\Psi_0(x)$ the digamma function. And finally the effective 4d Yukawa coupling constant turn out to be proportional to the physical mass according to

$$y_{4d} \approx \frac{1}{\sqrt{2}} \Gamma\left(\frac{3}{2} - \alpha\right) \left[\Psi_0\left(\frac{3}{2} - \alpha\right) + \ln\left(\frac{\tau_0}{\zeta_0}\right) \right] mk.$$

C Some technical aspects concerning the solution to the fermionic equations of motion

The system we are concerned with is of the form

$$D_1 h = g, \quad (116)$$

$$D_2 g = h, \quad (117)$$

where $D_{1,2}$ are some (linear) differential operators. The second order (decoupled) system is

$$D_2 D_1 h = h, \quad (118)$$

$$D_1 D_2 g = g. \quad (119)$$

Notice that whenever (116) and (118) hold then (117) and (119) are automatically satisfied and vice versa. In addition, let $\mathcal{H} = \{h : \mathbb{R} \rightarrow \mathbb{R} \mid D_2 D_1 h = h\}$ and $\Gamma = \{g : \mathbb{R} \rightarrow \mathbb{R} \mid D_1 D_2 g = g\}$ be the set of solutions to the second order differential equations (118) and (119), respectively. Then $\forall h \in \mathcal{H}$ ($\forall g \in \Gamma$), $D_1 h \in \Gamma$ ($D_2 g \in \mathcal{H}$). These considerations are useful when solving the first order constraints.

Concerning the orthonormality of the solutions to the system (26)-(27) we will show that the operators (28)-(29) are indeed hermitian with respect to the usual scalar product¹⁰, say

$$\langle \phi, \varphi \rangle = \int_{z_0}^{\infty} \phi^\dagger \varphi.$$

To see this notice that the second order differential operators can be compactly written as

$$\mathcal{O}H \equiv H'' - \left[f^2 (M + m_D \sigma_3)^2 - f' (M + m_D \sigma_3) - f m'_D \sigma_3 \right] H, \quad (120)$$

with $H = (h_+, h_-)^T$, and analogous expression for $G = (g_+, g_-)^T$. Then by means of partial integration we find

$$\langle \tilde{H}, \mathcal{O}H \rangle = \langle \mathcal{O}\tilde{H}, H \rangle - \tilde{H}^\dagger H' + \tilde{H}'^\dagger H \Big|_{z_0}, \quad (121)$$

and using the first order constraints

$$H' = - (M + m_D \sigma_3) f H - m G, \quad \tilde{H}' = - (M + m_D \sigma_3) f \tilde{H} - m \tilde{G},$$

¹⁰Whenever this scalar product be well defined, which is not the case for the non normalizable solutions.

the boundary piece reduces to

$$m \tilde{H}^\dagger G - m \tilde{G}^\dagger H. \quad (122)$$

Finally, the boundary conditions (30) allow us to deduce that $\tilde{H}^\dagger G = \tilde{H}^\dagger \sigma_1 (-\sigma_1) G = -\tilde{H}^\dagger G$ and thus $\langle \tilde{H}, \mathcal{O}H \rangle = \langle \mathcal{O}\tilde{H}, H \rangle$ for all $\tilde{H}, H$ solving the system (26)-(27). The same argument leads to the same conclusion for the operators (29).

D Fermionic zero-modes profile

Here we will plot the profiles of the wave functions corresponding to the top quark, for which $\alpha \approx 13$ and $\delta \sim 10^{-16}$




Figure 4: Plots for the profiles (from left to right and from top to bottom) h_+ (blue line), g_+ (red line), h_- (green line) and g_- (brown line) as a function of ζ for $\alpha = 13$ and $m k \sim 10^{-17}$, corresponding to the top quark in the Planckian case.

As can be seen the profiles are peaked towards the UV brane as opposed to the case of the hard wall scenario (RS) where the top is peaked in the IR brane. For the case of lighter fermions the behavior is qualitatively similar in both scenarios.

References

- [1] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999) [arXiv:hep-th/9906064];
L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221].
- [2] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **436**, 257 (1998) [arXiv:hep-ph/9804398]; N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **429**, 263 (1998) [arXiv:hep-ph/9803315].
- [3] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200]; S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109]; E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].
- [4] N. Arkani-Hamed, M. Porrati and L. Randall, JHEP **0108**, 017 (2001) [arXiv:hep-th/0012148]; R. Rattazzi and A. Zaffaroni, JHEP **0104**, 021 (2001) [arXiv:hep-th/0012248]; M. Perez-Victoria, JHEP **0105**, 064 (2001) [arXiv:hep-th/0105048].
- [5] A. Karch, E. Katz, D. T. Son and M. A. Stephanov, Phys. Rev. D **74**, 015005 (2006) [arXiv:hep-ph/0602229].
- [6] A. Falkowski and M. Perez-Victoria, JHEP **0812**, 107 (2008) [arXiv:0806.1737 [hep-ph]].
- [7] B. Batell, T. Gherghetta and D. Sword, Phys. Rev. D **78**, 116011 (2008) [arXiv:0808.3977 [hep-ph]].
- [8] C. Csaki and M. Reece, JHEP **0705**, 062 (2007) [arXiv:hep-ph/0608266]; U. Gurssoy and E. Kiritsis, JHEP **0802**, 032 (2008) [arXiv:0707.1324 [hep-th]]; U. Gurssoy, E. Kiritsis and F. Nitti, JHEP **0802**, 019 (2008) [arXiv:0707.1349 [hep-th]]; B. Batell and T. Gherghetta, Phys. Rev. D **78**, 026002 (2008) [arXiv:0801.4383 [hep-ph]].
- [9] F. Coradeschi, S. De Curtis, D. Dominici and J. R. Pelaez, JHEP **0804**, 048 (2008) [arXiv:0712.0537 [hep-th]]; C. P. Burgess, C. de Rham and L. van Nierop, JHEP **0808**, 061 (2008) [arXiv:0802.4221 [hep-ph]].
- [10] I.S. Gradshteyn and I.M. Ryzhik, "Table of Integrals, Seires and Products," American Journal of Physics (1988) Volume 56, Issue 10, pp. 958;
Herbert Buchholz, "The confluent Hypergeometric Function," Springer-Verlag (1969).